



**College of Agriculture and Environmental Sciences
 Department of Entomology
 Thrips as Vectors in Agricultural Commodities Lab**

The following report includes data obtained during the period February 7 – February 14 from Brooks, Colquitt, Decatur, and Tift Counties. Two fields were sampled in each of the 4 counties. Five of the target weeds were collected in one or more counties and included Carolina geranium, wild mustard, cudweed, sowthistle and chickweed. Thrips were extracted from the weeds using Berlese funnels, and the total number of thrips per county ranged from 11 to 78. ELISA was performed on 147 weed samples with 2.04 % indicating positive for tomato spotted wilt virus.

County and Field	Number of weed samples collected	Percent TSWV in weed samples	Total number of thrips collected	Number of <i>F. fusca</i>	Number of <i>F. occidentalis</i>
Brooks L1	6	0%	11	0	0
Brooks L2	6	5.6%	4	0	0
Colquitt L1	6	5.6%	17	0	0
Colquitt L2	6	0%	0	0	0
Decatur L1	6	0%	2	0	0
Decatur L2	7	0%	9	1	0
Tift L1	6	5.6%	5	0	0
Tift L2	6	0%	73	0	0

* Based on previous research, incidence of TSWV in more than 2% of weeds results in a high incidence year for the crop.

Stan Diffie
 Research Professional
 TVAC Lab
diffie@uga.edu

Dr. David Riley
 Associate Professor
 Dept. of Entomology
dgr@tifon.uga.edu